

# Health Outcomes & Patient Evaluation in Bipolar Disorder (HOPE.BD): A Canadian Prospective Study

P. Cervantes<sup>1</sup>, S. Beaulieu<sup>2</sup>, M. Connolly<sup>3</sup>, L. Yatham<sup>4</sup>, S. Ferrazzi<sup>5</sup>

<sup>1</sup>McGill University Health Centre, Montreal General Hospital Site, Montreal, QC; <sup>2</sup>Douglas Hospital & Research Center, McGill University, Montreal, QC; <sup>3</sup>Royal Jubilee Hospital, Victoria, BC; <sup>4</sup>University of British Columbia, UBC Hospital, Vancouver, BC; <sup>5</sup>Janssen-Ortho Inc., Toronto, Canada

## BACKGROUND<sup>1</sup>

- Bipolar Disorder (BD) is generally an episodic, lifelong illness with a variable course. The first episode of BD may be manic, hypomanic, mixed, or depressive. According to DSM-IV criteria, patients with BD Type I have had at least one episode of mania. Patients meeting criteria for BD Type II have a history of major depressive episodes and hypomanic episodes only.
- Bipolar Type I disorder affects approximately 0.8% of the adult population, with estimates from community sampling ranging from between 0.45% and 1.6%, affecting men and women fairly equally. Bipolar Type II disorder affects approximately 0.5% of the population and is apparently more common in women. These estimates of prevalence are considered conservative.
- Bipolar disorder causes substantial psychosocial morbidity, frequently affecting patients' relationships with spouses and partners, children, family members as well as occupation. It is estimated that as many as 60% of people diagnosed with BD-I experience chronic interpersonal or occupational difficulties and subclinical symptoms between acute episodes.

## RATIONALE

- There are many unanswered questions surrounding the human, social and economic costs of BD in Canada. Research is needed to establish the demographic, diagnostic and treatment factors associated with important outcomes such as euthymia, behavioural decline, suicide and health resource utilization.
- This study prospectively examines treatment patterns in bipolar disorder across Canada, for those patients under the care of a psychiatrist. Regional variations will be explored when and where appropriate.
- The focus of this publication is to present initial demographic and subject history.

## METHOD

- A prospective cohort study design involving a combination of University and Community hospitals/clinics from across Canada is currently enrolling 200 patients per study year with a target goal of 1000 patients over a 4-5 year period.
- Patients will be managed under conditions of routine clinical psychiatric practice in the context of accepted Canadian guidelines.
- With the patients' consent, data will be collected for at least a one-year follow-up period and entered into a computer database designed for psychiatric disorders.
- Community and health resource utilization will be collected through a home diary completed monthly by the patient.

## RESULTS

- The study began recruiting in April 2002 and currently has 14 active sites. As of March 2003, there were 145 patients enrolled from which data is available for 126.
- Mean age at study entry is 43.5 years (SD 13.5, range 18-80 yrs.) for the 75 females and 44.2 years (SD 11.9, range 20-72 yrs.), for the 54

males. Caucasians constitute 92% of the population, Asians 3% and Blacks 1%.

- Household income of less than \$40k was indicated by 54% of subjects, 24% had \$40k - \$69.9k, 8% between \$70 - \$99.9k and 9% above \$100k.
- Half of the subjects have children (49% BD-I & 55% BD-II) but only 24% reported that the children lived with them (8.7% unknown).
- Table 1 outlines basic demographic details for the subjects recruited thus far. Almost 3/4 have greater than high school education but appear to suffer from a high unemployment rate (22%). Approximately 25% have broken marriages.
- Although suicidality is higher in BD Type II (40.9%), it is not statistically different from the 28.0% in Type I (Figure 1).
- The proportion of subjects reporting prior substance abuse is relatively low in this population with overall rates ranging from 4% for opiates to 23% for alcohol (Table 2).
- A family history of mental health illness is the only parameter that demonstrates an association with bipolar disorder ( $p = 0.071$ , Figure 2).
- Hypomanic and depressive episodes are significantly ( $p < 0.05$ ) more prevalent in BD-II versus BD-I (Table 3).
- Mean patient age at the time of professional diagnosis was lower in Type I versus Type II (35 vs. 41 yrs.,  $p < 0.05$ ) however the mean age at time of first psychiatric admission was sooner in Type II disorder (18 vs. 29 yrs.,  $p < 0.05$ , Table 4).

Table 1. Demographics by Bipolar Type

Parameter	Bipolar I		Bipolar II		ALL	
	N	%	N	%	N	%
<b>Sex</b>						
Female	45	54.9	26	59.1	71	56.3
Male	37	45.1	18	40.9	55	43.7
<b>Marital Status</b>						
Unmarried	26	31.7	16	36.4	42	33.3
Married	18	22	17	38.6	35	27.8
Living together	11	13.4	4	9.1	15	11.9
Divorced	18	22	4	9.1	22	17.5
Separated	6	7.3	3	6.8	9	7.1
Widowed	1	1.2	na	na	1	0.8
Unknown	2	2.4	na	na	2	1.6
<b>Education</b>						
Less than High School	11	13.4	9	20.5	20	15.9
High School Graduate	7	8.5	4	9.1	11	8.7
College	20	24.4	9	20.5	29	23
Some University	15	18.3	7	15.9	22	17.5
Univ. Grad or post-graduate	26	31.7	15	34.1	41	32.5
Unknown	3	3.7	na	na	3	2.4
<b>Working Status</b>						
Full-time	26	31.7	13	29.5	39	31
Part-time	12	14.6	5	11.4	17	13.5
Housewife/-man	na	na	2	4.5	2	1.6
Unemployed	18	22	10	22.7	28	22.2
Studying	5	6.1	2	4.5	7	5.6
Retired	9	11	3	6.8	12	9.5
Other/Unknown	12	14.6	9	20.5	21	16.7

Figure 1. History of Suicidality




Figure 2. Family History




Table 2. Prior Substance Use/Abuse by Bipolar Disorder Type

Substance	Bipolar I		Bipolar II		ALL	
	N	%	N	%	N	%
<b>Alcohol</b>						
No	17	20.7	9	20.5	26	20.6
Use	27	32.9	18	40.9	45	35.7
Abuse	18	22	11	25	29	23
Unknown	20	24.4	6	13.6	26	20.6
<b>Stimulants</b>						
No	47	57.3	27	61.4	74	58.7
Use	10	12.2	8	18.2	18	14.3
Abuse	5	6.1	3	6.8	8	6.3
Unknown	20	24.4	6	13.6	26	20.6
<b>Cannabis</b>						
No	28	34.1	22	50	50	39.7
Use	23	28	11	25	34	27
Abuse	11	13.4	5	11.4	16	12.7
Unknown	20	24.4	6	13.6	26	20.6
<b>Opiates</b>						
No	53	64.6	32	72.7	85	67.5
Use	5	6.1	5	11.4	10	7.9
Abuse	4	4.9	1	2.3	5	4
Unknown	20	24.4	6	13.6	26	20.6

Table 3. Onset and Course of Illness

	Bipolar I			Bipolar II			ALL		
	N	Mean	STD	N	Mean	STD	N	Mean	STD
Hospital Admissions for BD	62	1.2	1.4	38	0.5	1.1	100	0.9	1.3
Hypomanic Episodes	62	1.9	6.0	37	6.7	10.9	99	3.7	8.5
Manic Episodes	62	0.7	1.2	38	0	0.16	100	0.5	1.0
Depressive Episodes	62	2.2	2.9	37	6.2	10.6	99	3.7	7.1
Psychotic Episodes	62	0.1	0.4	38	0.1	0.32	100	0.1	0.3

Table 4. History of Illness by Bipolar Disorder Type

	Bipolar I			Bipolar II			ALL		
	N	Mean	STD	N	Mean	STD	N	Mean	STD
At what age was professional help first sought for bipolar disorder symptoms?	62	30	13.5	38	31	11.4	100	30	12.7
At what age was there a professional diagnosis of bipolar disorder?	62	35	12.4	38	41	10.9	100	37	12.2
At what age did the subject first try a mood stabilizing (bipolar) medication?	62	34	13	38	38	13	100	36	13
At what age was the subject's first psychiatric admission?	62	29	16.4	37	18	19.9	99	25	18.4

## CONCLUSIONS

- This is the first Canada wide, prospective observational study to rigorously examine bipolar disorder.
- The burden of bipolar disorder on society and patients is demonstrated by the early age of onset, high levels of unemployment/suicide/substance abuse and the chronic nature of the disease.
- This "practice based" approach is a useful tool for exploring relationships between relevant patient types, and will provide a means of elucidating the effectiveness of different treatment strategies and economic parameters in BD in Canada.

## REFERENCES

- Supplement to The American Journal of Psychiatry. Practice Guideline for the Treatment of Patients with Bipolar Disorder (Revision). Volume 159, Number 4, April 2002.

## OTHER STUDY SITES

- Balaraju Katta/Terry Isamura/Hem Phaterpekar, Royal Columbian Hospital, New Westminster, BC; Javed Ali, Cape Breton Regional Hospital, Sydney, NS; Chris Gorman, Foothills Hospital, Calgary, AB; Peter Silverstone, Edmonton, AB; Verinder Sharma, Regional Mental Health Care, London, ON; Mark Lander, Winnipeg, Man; Sagor Parikh, CAMH, Toronto, Ont; Philippe Baruch, Centre Hospitalier Robert-Gifford, QC; Jean Leblanc, Hôpital du Sacre-Coeur de Montreal, QC; Roumen Milev, Kingston, ON